



MONOCYTE CHEMOATTRACTANT ACTIVITY OF GALECTIN-3

RECEIVED
FEB 28 2002
TECH CENTER 1600/2900

Statement as to Federally Sponsored Research

This work was supported in part by National Institute of Health Grant No. RO1

5 AI39620. The government may have certain rights in the invention.

Cross Reference to Related Applications

This application is based on Provisional Application Ser. No. 60/188,795, which was
filed on Mar. 13, 2000, and priority is claimed thereto.

10

Field of the Invention

The present invention relates to methods for modulating migration of cells, especially
monocytes, neutrophils and macrophages, using galectin-3, galectin-3 binding polypeptides,
galectin-3 receptor binding polypeptides or galectin-3 mimetics. The invention also relates to
15 screening methods for identifying agents that modulate galectin-3-mediated cell migration.

Background of the Invention

Lectins are proteins that bind to specific carbohydrate structures and can thus
recognize particular glycoconjugates. Galectins are a family of over 10 structurally related
20 lectins that bind beta-galactosides.

Galectin-3 is a 26 kDa beta-galactoside-binding protein belonging to the galectin
family. This protein is composed of a carboxyl-terminal carbohydrate-recognition domain
(CRD) and amino-terminal tandem repeats. Galectin-3 is found in epithelia of many organs,
as well as in various inflammatory cells, including macrophages, dendritic cells and Kupffer
25 cells. The expression of galectin-3 is upregulated during inflammation, cell proliferation, cell
differentiation, and through transactivation by viral proteins. Its expression is also affected by
neoplastic transformation -- upregulated in certain types of lymphomas and thyroid
carcinoma; downregulated in other types of malignancies, such as colon, breast, ovarian and
uterine carcinomas. Recently, it has been reported that the expression of this lectin has a
30 strong correlation with the grade and malignant potential of primary brain tumors. Increased
galectin-3 expression has also been noted in human atherosclerotic lesions. These findings
suggest that galectin-3 may mediate both physiological and pathological responses.